

Bersanlimab Biosimilar - Research Grade

Product Code: ICH5394

More Information

Species Reactivity: Human

Target: ICAM1 $\frac{1}{2}$

Concentration: 5mg/ml

Isotype: IgG1 - lambda2

Host: CHO Cells

product name H2: Bersanlimab Biosimilar - Research Grade

Sequence: <https://www.imgt.org/mAb-DB/>

Buffer: ICH3002

UniProt: P05362;P13597

Shipping Conditions: Blue ice

background: Intercellular adhesion molecule 1 (ICAM1/CD54) mediates leukocyte adhesion and transmigration and is a recurring target in inflammation and vascular-biology research. This product is an unconjugated, non-therapeutic analog of bersanlimab, a human IgG1-lambda recombinant antibody supplied for research use only. It is an ultra-pure, low-endotoxin preparation built to in vivo standards and produced in scalable milligram-to-gram quantities for bulk and high-throughput laboratories. Applications include functional and adhesion assays, leukocyte-trafficking studies, in vitro screening and preclinical models that interrogate ICAM-1-dependent inflammatory processes. As a research-use-only reagent, it combines a low-endotoxin in vivo-grade format with scalable milligram-to-gram production for demanding studies.

Purification Method: This monoclonal antibody was purified using Protein A

Formulation: Sterile, preservative-free, solution in PBS. BSA and Azide free.

ichor.bio // hello@ichor.bio

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

ichorbio, ichorbio logo and all other trademarks are the property of ichorbio LTD © ichorbio

2026



Purity: >90% by SDS-PAGE

Endotoxin: ≤ 1.0 EU/mg as determined by the LAL method

Storage: This biosimilar is stable when stored at 2-8°C. For longer term storage (> 3 months) it is recommended to store this antibody at -20°C or -70°C.

Applications: Functional Assays

Application Notes: This biosimilar is for research use only (RUO): it is not for diagnostic or therapeutic procedures and cannot be purchased by patients.

Use: Products are for research use only. Not for use in diagnostic or therapeutic procedures.

Delivery Time: 3-4 weeks