

Human CD16a (176F) Protein (In Vivo Grade, Ultra-Low Endotoxin)

Product Code: ICH3014

More Information

Species Reactivity: Human

Target: Human CD16a (176F) Protein

Host: Human embryonic kidney (HEK) 293 cells.

product name H2: Human CD16a (176F) Protein (In Vivo Grade, Ultra-Low Endotoxin)

UniProt: P08637

Shipping Conditions: Ambient

background: The 176F (low-affinity) allotype of human CD16a (FcγRIIIa) is a major activating Fc gamma receptor on NK cells and macrophages that drives antibody-dependent cellular cytotoxicity, and allelic variation at this position strongly influences therapeutic-antibody potency. This recombinant CD16a (176F) ectodomain, expressed in HEK 293 cells, is validated for SPR, BLI and ELISA to support Fc binding kinetics, ADCC characterization and Fc-variant ranking. It is a research-use-only protein supplied in an in vivo grade, ultra-low-endotoxin format and produced at scalable milligram (mg) to gram (g) scale, giving high-throughput and bulk-screening laboratories a dependable allotype-defined CD16a reagent.

Other names: Human Fc gamma RIIIa, CD16A, FCGR3A, FCGRIIIA, FCR3A, FCRIIIA, IGFR3A, IGFRIIIA

Specificity: The sequence of the extracellular domain of human CD16a (Gly 17-Gln 208) was fused with a C-terminal tag consisting of the AVI tag, TEV protease recognition sequence and a 10-His tag.

Formulation: Lyophilized from sterile PBS, pH 7.4. No preservatives or cryoprotectants have been added. To obtain a final concentration of 1 mg/ml reconstitute 0.25 mg vials with



0.25 ml water and 1.0 mg vials with 1.0 ml water.

Purity: >95% monomer purity as determined by SDS-PAGE and SEC-HPLC.

Endotoxin: 1.0 EU per mg as determined by the LAL method.

Storage: Lyophilized proteins are stable at ambient temperature for at least 2 weeks. If the protein is not to be used immediately then the protein should be stored in lyophilized form at -20 °C for up 12 months. Once the protein has been reconstituted we recommen

Applications: SPR, BLI, ELISA

Use: Products are for research use only. Not for use in diagnostic or therapeutic procedures.